




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07D 487/04, A61K 31/5025 // (C07D 487/04, 237:00, 257:00) (C07D 487/04, 237:00, 249:00)	A1	(11) International Publication Number: WO 00/26218 (43) International Publication Date: 11 May 2000 (11.05.00)
(21) International Application Number: PCT/EP99/07304 (22) International Filing Date: 1 October 1999 (01.10.99) (30) Priority Data: MI98A002319 29 October 1998 (29.10.98) IT (71) Applicant (for all designated States except US): ZAMBON GROUP S.P.A. [IT/IT]; via della Chimica, 9, I-36100 Vicenza (IT). (72) Inventors; and (75) Inventors/Applicants (for US only): NAPOLETANO, Mauro [IT/IT]; via Venini, 37, I-20127 Milano (IT). NORCINI, Gabriele [IT/IT]; via A. Volta, 42, I-21010 Vizzola Ticino (IT). PELLACINI, Franco [IT/IT]; via G. Balla, 14, I-20151 Milano (IT). MORAZZONI, Gabriele [IT/IT]; via Labriola, 12, I-20020 Lainate (IT). (74) Agent: LONGONI, Alessandra; Zambon Group S.p.A., Corp. Patent & Trademark Dept., via Lillo del Duca, 10, I-20091 Bresso (IT).		(81) Designated States: AU, CA, CZ, HU, IL, JP, KR, NZ, SI, US, ZA, Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>

(54) Title: TRICYCLIC PHTHALAZINE DERIVATIVES AS PHOSPHODIESTERASE 4 INHIBITORS**(57) Abstract**

Tricyclic phthalazine compounds of formula (I) wherein A is a 5-7 membered heterocycle containing from 1 to 4 nitrogen atoms, optionally partially or totally unsaturated, and optionally substituted by a (C₁₋₄)alkyl group in turn optionally substituted; Z is NH, methylene, a C₂₋₆ alkylene chain optionally branched and/or unsaturated and/or interrupted by a C₅₋₇ cycloalkyl residue; Cy is phenyl or heterocycle optionally substituted by one or more substituents, or a COR₄ group wherein R₄ is hydroxy, alkoxy, amino optionally substituted by one or two (C₁₋₆)alkyl groups or by hydroxy; R is a (C₁₋₆)alkyl or polyfluoro(C₁₋₆)alkyl group; R₁ is hydrogen; a (C₁₋₈)alkyl, (C₂₋₈)alkenyl or (C₂₋₈)alkynyl group optionally substituted by hydroxy, oxo, aryl or heterocycle, and optionally interrupted by one or more heteroatoms or heterogroups; a (C₁₋₄)alkoxy group or a (C₄₋₇)cycloalkoxy group optionally containing an oxygen atom and optionally substituted by a polar substituent in the cyclic moiety, aryloxy, aryl-(C₁₋₁₀)-alkoxy; the N-O derivatives and the pharmaceutically acceptable salts thereof are described. The compounds of formula (I) are PDE 4 inhibitors.



BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

TRICYCLIC PHTHALAZINE DERIVATIVES AS PHOSPHODIESTERASE 4 INHIBITORS

The present invention relates to tricyclic derivatives, to the pharmaceutical compositions
5 containing them and to their use as phosphodiesterase 4 inhibitors.


Phosphodiesterases are a family of isoenzymes which constitute the basis of the main
mechanism of cAMP (cyclic adenosine-3',5'-monophosphate) hydrolytic inactivation.
cAMP has been shown to be the second messenger mediating the biologic response to many
of hormones, neurotransmitters and drugs [Krebs Endocrinology Proceedings of the 4th
10 International Congress Excerpta Medica, 17-29, 1973]. When the suitable agonist binds to
the cell surface, the adenylated cyclase activates and turns Mg^{2+} -ATP into cAMP. cAMP
modulates the activity of the majority, if not of all the cells contributing to the
pathophysiology of various respiratory diseases, both of allergic origin and not. It follows
that an increase of the cAMP concentration yields beneficial effects such as airway smooth
15 muscle relaxation, inhibition of the mast cell mediator release (basophil granulose cells),
suppression of the neutrophil and basophil degranulation, inhibition of the monocyte and
macrophage activation. Thus, compounds able of activating adenylate cyclase or of
inhibiting phosphodiesterases could suppress the undesired activation of the airway smooth
muscle and of a great number of inflammatory cells.

20 In the phosphodiesterase family there is a distinct group of isoenzymes, phosphodiesterases 4
(hereinafter PDE 4) specific for the cAMP hydrolysis in the airway smooth muscle and
inflammatory cells (Torphy, "Phosphodiesterase Isoenzymes: Potential Targets for Novel
Anti-asthmatic Agents" in New Drugs for Asthma, Barnes, ed. IBC Technical Services Ltd,
1989). Studies carried out on this enzyme show that its inhibition yields not only the airway
25 smooth muscle relaxation, but also the suppression of mastocyte, basophil and neutrophil
degranulation, so as the inhibition of the monocyte and neutrophil activation. In addition, the
action of PDE 4 inhibitors is markedly strengthened when the adenylate cyclase activity of
the target cells is increased by endogenous hormones, as it happens *in vivo*. Thus PDE 4
inhibitors are effective in the therapy of asthma. Such compounds offer a unique approach to
30 the therapy of various respiratory diseases, both of allergic origin and not, and possess
significant therapeutic advantages over the current therapy.

- 2 -


The excessive or irregular production of tumour necrosis factor (hereinafter TNF_α), a cytokine with pro-inflammatory activity produced by various kinds of cells, affects the mediation or the exacerbation of many pathologies such as, for example, the adult respiratory distress syndrome (ARDS) and the chronic pulmonary inflammatory disease. Therefore compounds able to control the negative effects of TNF_α , i.e. the inhibitors of this cytokine, are to be considered as useful against many pathologies.

The patent EP 0 526 840 (iKyowa Hakko Kogyo) claims compounds of formula



wherein R_1 is hydrogen, (C_{1-6}) alkyl, (C_{7-15}) arylalkyl or optionally substituted aryl; and R_2 is hydrogen, (C_{1-6}) alkyl, thienyl or optionally substituted aryl. These compounds are said to be active, *inter alia*, as antiinflammatories, immunosuppressives, bronchodilators.


The patent application JP 09227563 (Lederle Japan) describes compounds of formula



wherein R is an optionally substituted amino group, Z is S or O , A and B form a benzene ring or are absent, and n is 0-2. These compounds are useful as bronchodilators, antiasthmatics, antihypertensives and anticholesterolemics.

The patent application WO 97/34893 (Astra) describes compounds of formula

- 3 -




5

wherein B, D, E and G may form a benzene ring optionally substituted by alkoxy; X is C=O, C=S, C=NR, CR₃R₆ or NR₄; R₃ is H or forms a bond with R₂; R₄ is lower alkyl or forms a bond with R₂; R₆ may be H, lower alkyl optionally substituted by phenyl, or cycloalkyl, phenyl, etc.; Y is N or CR; R₂ may be H, lower alkyl optionally substituted by phenyl, COOR, NR'R'', OR, F, or cycloalkyl or may form a bond with one of R₁, R₃ and R₄; R₁ may be OH or lower alkyl or may form a bond with one of R₂ and R₅; R₅ is a bond with R₁ or R₆; Z is OR₈ or O; and Ar₁ may be optionally substituted phenyl, pyridyl, pyrimidyl. These compounds have an antiinflammatory activity.

15

The patent application WO 98/09969 (Astra) describes compounds of formula



20


wherein A, A₁, A₂ and A₃ may be CH or CR₄; X may be CH₂ or O; Y may be a bond, CH₂, C=O, C substituted by alkyl in turn substituted by a cyclic residue; Z is a bond or CH₂; R₁ is hydrogen, lower alkyl or alkoxy; R₂ and R₃ are hydrogen or form a bond; and R₄ may be optionally substituted alkoxy. These compounds have an antiinflammatory and antiallergic activity.

25

The patent application DE 19617862 (Schering AG) describes compounds of formula

30


- 4 -



5

10 wherein, *inter alia*, R_1 and R_2 are H, alkyl, nitro, halogen, amino, lower alkoxy, CF_3 ; R_3 and R_4 are H, alkyl, aryl, heteroaryl or cycloalkyl; $X=H$; Y is alkoxy or $X+Y = -O-(CH_2)_n-O-$; $n=1-3$; and A is a 5-member heterocycle having from 1-3 nitrogen atoms. These compounds are inhibitors of glutamate receptor.

The patent application EP 0 548 923 (Takeda Chemical Ind.) describes, *inter alia*,
 15 compounds of formula




20


25 wherein R_1 is H or a lower alkyl group or a halogen atom; R_4 and R_5 are H or a lower alkyl group or form a 3-7 membered cycle optionally containing a heteroatom together with the carbon atom to which they are bound; A is an optionally substituted amino group; and $m, n = 1-4$. These compounds are antiallergics, antiinflammatories and anti-PAF (anti-piastrinic activating factor), and are useful as antiasthmatics. In fact, these compounds act through an anti-PAF mechanism which makes them bronchodilators.

Similar compounds are claimed in the patent application EP 0 620 224 (Takeda Chemical
 30 Ind.) which illustrates, *inter alia*, the general formula

- 5 -




wherein R_1 is H or a lower alkyl group or a halogen atom; X is an oxygen or sulphur atom or a $-CH_2-$ group; Y is a group



wherein R_4 and R_5 are H or a lower alkyl group, or is a 3-7 membered cycle optionally containing a heteroatom; R_6 and R_7 are H, an optionally substituted lower alkyl, cycloalkyl or aryl or together with the nitrogen atom to which they are bound form a heterocycle; and $m, n=0-4$. These compounds have the same activity claimed in the just above cited patent application.

The patent application WO 98/21208 (Byk Gulden Lomberg) claims PDE3 and PDE4 inhibitors of formula



wherein, *inter alia*, R_1 is an alkyl group; R_2 and R_3 are hydroxy, optionally fluorinated alkoxy, cycloalkoxy and cycloalkylmethoxy; and R_4 is a phenyl group substituted by carboxy, amido or alkoxycarbonyl and optionally substituted by halogen, alkyl, CF_3 , nitro or hydroxy. These compounds are said to be useful in the treatment of pathologies of the airways and/or of dermatosis.


It has now surprisingly been found a new class of phthalazine derivatives able to inhibit PDE

- 6 -

4 and TNF_α .

Therefore the present invention relates to compounds of formula

5



wherein

- 10 A is a 5-7 membered heterocycle containing from 1 to 4 nitrogen atoms, optionally partially or totally unsaturated, and optionally substituted by a (C_{1-4}) alkyl group in turn optionally substituted;

Z is NH , methylene, a C_{2-6} alkylene chain optionally branched and/or unsaturated and/or interrupted by a C_{5-7} cycloalkyl residue;

- 15 Cy is phenyl or heterocycle optionally substituted by one or more substituents, or a COR_4 group wherein R_4 is hydroxy, alkoxy, amino optionally substituted by one or two (C_{1-6}) alkyl groups or by hydroxy;

R is a (C_{1-6}) alkyl or polyfluoro (C_{1-6}) alkyl group:

- R_1 is hydrogen; a (C_{1-8}) -alkyl, (C_{2-8}) -alkenyl or (C_{2-8}) -alkynyl group optionally substituted by
20 hydroxy, oxo, aryl or heterocycle, and optionally interrupted by one or more heteroatoms or heterogroups; a (C_{1-4}) alkoxy group or a (C_{4-7}) cycloalkoxy group optionally containing an oxygen atom and optionally substituted by a polar substituent in the cyclic moiety, aryloxy, aryl- (C_{1-10}) -alkoxy;

- the $\text{N} \rightarrow \text{O}$ derivatives of the compounds of formula I and the pharmaceutically acceptable
25 salts thereof.

Preferred compounds according to the invention are those of formula I wherein Z is methylene or a C_{2-6} alkylene chain.

- Still more preferred compounds according to the invention are those of formula I wherein Z is methylene or a C_{2-6} alkylene chain; and Cy is a heterocycle optionally substituted by one
30 or more substituents.

- 7 -

Still more preferred compounds according to the invention are those of formula I wherein Z is methylene; and Cy is pyridine substituted by two substituents.

The compounds of formula I can have one or more asymmetric centres and therefore they
5 can be in the form of stereoisomers. Object of the present invention are compounds of formula I in the form of diastereoisomeric mixtures as well as of single stereoisomers.

The compounds of formula I are active as PDE 4 and TNF_{α} inhibitors and thus are used as therapeutic agents in allergic and inflammatory pathologies such as, for example, emphysema, chronic bronchitis, asthma and allergic rhinitis.

10 For substituent Cy, as heterocycle it is particularly meant pyrrole, imidazole, pyrazole, pyridine, pyrazine, pyrimidine, pyridazine, piperazine, triazine, morpholine, pyrrolidine, pyrroline, imidazoline, pyrazoline, pyrazolidine, imidazolidine, piperidine, furan, pyran, isothiazole, isoxazole, thiophene and the like.

The optionally present substituents can be oxo, nitro, carboxy, halogen, that means a
15 fluorine, chlorine, bromine or iodine atom. As "polar substituent" they are meant those groups made by atoms having a different electronegativity, so forming a dipole, such as, for example, a hydroxy or keto group.

Specific examples of alkyl groups are methyl, ethyl, n-propyl, i-propyl, n-butyl, s-butyl, t-butyl, n-pentyl, 1-methyl-butyl, 2-ethyl-propyl, 3-methyl-butyl, 3-methyl-2-butyl, n-hexyl,
20 heptyl, octyl and the like; examples of substituents optionally present on the alkyl groups are (C_{1-6}) alkoxy groups and amino groups mono- or di-substituted by (C_{1-4}) alkyl groups.

As (C_{4-7}) cycloalkyl group cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl are meant, while the aryl and the aryl moiety of the aryl- (C_{1-10}) alkyl substituent mean an aromatic ring of 6-10 carbon atoms such as, for example, phenyl, naphthyl, indanyl, and the like, and,
25 consequently, as aryl- (C_{1-10}) -alkyl substituent, benzyl, phenylethyl, phenyl-pentyl, indanyl-pentyl and the like.


The oxidised form $N \rightarrow O$, when present, can be on the nitrogen atoms of the phthalazine ring as well as on those on Cy.

Pharmaceutically acceptable salts of the compounds of formula I are those with organic and
30 inorganic acids, such as, for example, hydrochloric, hydrobromic, hydroiodic, sulfuric,


- 8 -

phosphoric, nitric, acetic, benzoic, maleic, fumaric, succinic, tartaric, citric, aspartic, methanesulfonic and 3,7-di-t.butyl-naphthalen-1,5-disulfonic (dibudinic acid). or with inorganic bases such as, for example, sodium or potassium hydroxide, sodium bicarbonate.


- 5 The synthesis of the compounds of formula I proceeds according to methods known to the skilled in the art. For example, when a compound of formula I wherein Z is different from NH is desired, the synthesis can start from an acid of formula




wherein R and R₁ are as above defined, which by reaction with formaldehyde/HCl gives a compound of formula



wherein R and R₁ are as above defined. This is oxidised, for example, with benzoyl peroxide/N-bromo-succinimide, and then hydrolysed to give a compound of formula




wherein R and R₁ are as above defined, which is treated with a hydrohalogenidric acid (HX) and triphenylphosphine to give a compound of formula



- 9 -

wherein R and R₁ are as above defined. This compound can be also obtained from compound III by radicalic halogenation with, for example, azaisobutyronitrile or benzoyl peroxide/N-bromo- or chloro-succinimide to give the compound of formula

5




10 wherein R and R₁ are as above defined, and X is chlorine or bromine, which gives compound V by treatment with triphenylphosphine.

Compound V is treated with an aldehyde of formula




15 wherein Cy is as above defined and Z'' is a C₁₋₅ alkylene chain optionally branched and/or unsaturated and/or interrupted by a C₅₋₇ cycloalkyl residue, or it is absent, in the presence of an organic base such as, for example, triethylamine, and gives a compound of formula

20



wherein R, R₁, Z'' and Cy are as above defined. This is reacted with hydrazine to give a compound of formula


25



30 wherein R, R₁, and Cy are as above defined and Z has the meanings reported in formula I but amino, which is treated with a halogenating agent, such as phosphoryl chloride or bromide,

- 10 -

to give a compound of formula



10 wherein R, R₁ and Cy are as above defined, X is chlorine or bromine, and Z is different from amino, which is treated with a suitable nucleophile such as, for example, sodium azide or sodium tetrazolate, or with hydrazine and then with a suitable acylating agent such as, for example, acetic anhydride or acetyl chloride, and gives the desired compound of formula I.

The synthesis of the N-oxides of the compounds of formula I occurs by treating the compounds of formula I with peracids such as, for example, m-chloroperbenzoic acid.

15 The preparation of the salts of the compounds of formula I is carried out according to conventional methods.

The compounds of formula I are PDE 4 inhibitors as showed by the *in vitro* enzymatic inhibition tests (example 18), and furthermore are able to inhibit the TNF_α release.

20 It is apparent how these enzymatic selectivity and specificity features combined with the lack of activity on the cardiovascular system make the compounds of formula I specifically suitable for treating pathologies involving PDE 4 and TNF_α even if in the present context the interest is particularly focused on the respiratory pathologies. In particular the compounds of the invention are useful for treating allergic and inflammatory diseases and above all for treating emphysema, chronic obstructive pulmonary disease (COPD) and chronic bronchitis

25 specifically, asthma and allergic rhinitis.

The therapeutic doses shall be generally from 0.1 to 1,000 mg a day and from 1 to 100 mg by oral route for single administration.

A further object of the present invention are the pharmaceutical compositions containing a therapeutically effective amount of the compounds of formula I or pharmaceutically

30 acceptable salts thereof in admixture with a suitable carrier.

- 11 -

The pharmaceutical compositions object of the invention can be liquid, suitable for the enteral or parenteral administration, and, preferably, solid such as tablets, capsules, granulates, suitable for the oral administration, or in a form suitable for the transdermal and
5 inhalatory administration.

The preparation of the pharmaceutical compositions object of the invention can be carried out according to common techniques.

In order to better illustrate the invention the following examples are now given.

The ¹H-NMR spectra were run at 200 MHz, δ are in parts per million.

10 Example 1

5,6-Dimethoxy-3H-isobenzofuran-1-one

A suspension of 3,4-dimethoxy-benzoic acid (353.5 g, 1.94 moles) in HCHO (1.7 l, 24.5 moles) was prepared under mechanic stirring, cooled in ice, saturated with gaseous HCl (340 g, 9.32 moles), then gradually brought to 60°C. After one night the temperature was brought
15 to the room values and further HCl (300 mg) was bubbled, then the temperature was brought again to 60°C for one night. The mixture was brought to small volume, taken up with water (1 l), neutralised with 28% NH₄OH (1.5 l) and kept at cool for 2 hours, then filtered. The filtrate was washed with water up to neutrality, then crystallised from CH₃OH (2 l) and dried under vacuum at 60°C to give 220 g of the title compound (yield: 58.65%).

20 ¹H-NMR (CDCl₃): 7.28 and 6.28 (2s, 2H); 5.20 (s, 2H); 3.95 and 3.91 (2s, 6H).

Example 2

2-Formyl-4,5-dimethoxy-benzoic acid

A mixture of 5,6-dimethoxy-3H-isobenzofuran-1-one (10 g, 51.5 mmoles), obtained as described in example 1, under N₂ in CCl₄ (250 ml), N-bromo-succinimide (13.88 g, 77.25
25 mmoles) and benzoyl peroxide (320 mg, 1.23 mmoles) was kept under reflux for 2 hours, then cooled, filtered and washed with a 10% Na₂SO₃ solution (200 ml), then with water, anhydried and brought to dryness. The residue was taken up with 5% HCl (100 ml) and kept under reflux for 4 hours, then the solution was cooled, basified with NaOH, washed with ethyl acetate and slowly re-acidified to give a precipitate which was filtered, washed
30 with water, dried on P₂O₅ under vacuum to give 6.43 g of the title compound (yield: 60%).

- 12 -

Example 3

5,6-Dimethoxy-3-(triphenyl- λ^6 -phosphanyl)-3H-isobenzofuran-1-one

A suspension of 2-formyl-4,5-dimethoxy-benzoic acid (6.43 g, 30.62 mmol), obtained as described in example 2, triphenyl-phosphine (8.3 g, 30.62 mmol), 30% HBr in acetic acid (8.26 ml, 30.62 mmol) and glacial acetic acid (20 ml) under N₂ was heated at 90°C for 4.5 hours. The mixture was brought to dryness, re-dissolved in acetonitrile (50 ml) and diluted with ethyl ether up to turbidity, then cooled and filtered, and the filtrate was washed with ethyl ether and dried under vacuum to give 13.6 g of the title compound (yield: 83%).

¹H-NMR (DMSO): 8.35 and 7.31 (2s, 2H); 8.03-7.66 (m, 15H); 6.01 (s, 1H); 3.84 and 3.45 (2s, 6H).

Example 4

5,6-Dimethoxy-3-pyridin-4-ylmethylen-3H-isobenzofuran-1-one

Triethylamine (20 ml, 145 mmol) was dropwise added to a suspension of 5,6-dimethoxy-3-(triphenyl- λ^6 -phosphanyl)-3H-isobenzofuran-1-one (78 g, 145 mmol), obtained as described in example 3, and 4-pyridincarboxaldehyde (13 ml, 145 mmol) in CH₂Cl₂ (1 l), at room temperature under stirring. After 1.5 hours the mixture was filtered and evaporated and the residue was treated with ethanol under reflux, cooled and filtered. The mother liquors were chromatographed (eluent: 100% CH₂Cl₂, then with 1% CH₃OH) and the residue was brought to dryness and joined to the above filtrate to give 25 g of the title compound.

Example 5

6,7-Dimethoxy-4-pyridin-4-ylmethyl-2H-phthalazin-1-one

5,6-Dimethoxy-3-pyridin-4-ylmethylen-3H-isobenzofuran-1-one (25 g, 88.34 mmol), obtained as described in example 4, was reacted with hydrazine hydrate (500 ml) for 2 hours at room temperature under stirring, then for 1 hour under reflux. The mixture was diluted with water (300 ml), cooled and filtered to give 23 g of the title compound (yield: 87%).

Example 6

1-Chloro-6,7-dimethoxy-4-pyridin-4-ylmethyl-phthalazine

A suspension of 6,7-dimethoxy-4-pyridin-4-ylmethyl-2H-phthalazin-1-one (10 g, 33.6 mmol), obtained as described in example 5, in POCl₃ (70 ml) was heated at 90°C for 4

- 13 -

hours. POCl_3 was evaporated and the residue dissolved in water, a saturated NaHCO_3 solution and NaOH up to obtain a precipitate which was filtered and re-suspended in CH_3OH , brought to dryness, re-suspended in acetone and filtered again. The residue was
5 dried at 45°C under vacuum to give 9.56 g of the title compound.

Example 7

8,9-Methoxy-6-pyridin-4-ylmethyl-tetrazol[5,1-a]phthalazine (Compound 1)

NaN_3 (103 mg, 1.583 mmoles) was added to a solution of 1-chloro-6,7-dimethoxy-4-pyridin-4-ylmethyl-phthalazine (500 mg, 1.583 mmoles), obtained as described in example 6, in
10 DMF (4.5 ml) and the mixture was heated at 80°C for 16 hours. DMF was evaporated and the residue partitioned between water and CH_2Cl_2 . The collected organic phases were anhydri-fied and brought to residue to give 420 mg of the title compound (yield: 82.3%).
 $^1\text{H-NMR}$ (CDCl_3): 8.55-8.52 (m, 2H); 8.00 (s, 1H); 7.25-7.22 (m, 3H); 4.59 (s, 2H); 4.10 and 3.90 (2s, 6H).

15

Example 8

6-Methoxy-3H-isobenzofuran-1-one

Formaldehyde 48% v/v (65 ml, 0.86 moles) under stirring, then 3-methoxy-benzoic acid (100 g, 0.66 moles) were added to concentrated HCl (1 l) and the mixture was heated at 100°C by checking the development of gas for 30 minutes. The cooling of the mixture
20 brought to the formation of a precipitate which was filtered and put aside, while the mixture was washed with water, then with 5% NaOH . The new precipitate was extracted twice with CH_2Cl_2 , the extract was anhydri-fied, concentrated, joined to the previously filtered solid, and both were dissolved in CH_2Cl_2 and treated with diethylamine (120 ml, 1.15 moles). After 24 hours it was extracted with 10% HCl and the phases were separated with CH_2Cl_2 . The
25 organic phase was washed with 10% NaOH , decoloured with charcoal, anhydri-fied and concentrated. The residue was dissolved in CH_2Cl_2 and treated, under stirring, with 10% HCl for 30 minutes. The organic phase was washed with water, anhydri-fied and concentrated. The residue was dissolved in CH_2Cl_2 and treated with 10% NaOH under stirring for 30 minutes. The organic phase was washed with water, dried and concentrated to give a solid
30 which was crystallised from aqueous CH_3OH . The filtrate was dried at 50°C on P_2O_5 , then

- 14 -

crystallised again from aqueous CH₃OH to give 35.28 g of the title compound (yield: 32%).

¹H-NMR (CDCl₃): 7.37-7.20 (m, 3H); 5.21 (s, 1H); 3.85 (s, 3H).

Example 9

5 3-Bromo-6-methoxy-3H-isobenzofuran-1-one

6-Methoxy-3H-isobenzofuran-1-one (35.28 g, 0.215 moles), obtained as described in example 8, suspended in CCl₄ (350 ml) under N₂, was added with N-bromo-succinimide (40 g, 0.225 moles), benzyl-peroxide in catalytic amount, then was slowly brought under reflux. After 2.5 hours the heating was stopped and the mixture was left standing overnight at room
10 temperature. Further catalyst was added and it was heated for further 3.5 hours. The mixture was cooled in ice and filtered over celite by washing well with CCl₄, then concentrated to give 41 g of the title compound (yield: 78%).

¹H-NMR (CDCl₃): 7.50-7.25 (m, 4H); 3.87 (s, 3H).

Example 10

15 (5-Methoxy-3-oxo-1,3-dihydro-isobenzofuran-1-yl)triphenylphosphonium bromide

Triphenylphosphine (42 g, 0.16 moles) was added to 3-bromo-6-methoxy-3H-isobenzofuran-1-one (41 g, 0.169 moles), obtained as described in example 9, suspended in anhydrous acetonitrile (205 ml) under N₂. The mixture was heated under reflux and after 3 hours cooled and concentrated to give a solid which was treated with ethyl ether, filtered and concentrated
20 under vacuum. There were thus obtained 74 g of the title compound (yield: 84%).

¹H-NMR (CDCl₃): 9.63 (s, 1H); 7.84-7.75 (m, 15H); 7.09-6.91 (m, 3H); 3.77 (s, 3H).

Example 11

3-(3,5-Dichloro-pyridin-4-ylmethylen)-6-methoxy-3H-isobenzofuran-1-one

Triethylamine (18.5 ml, 0.134 moles) was dropwise added to a suspension of (5-methoxy-3-oxo-1,3-dihydro-isobenzofuran-1-yl)triphenyl phosphonium bromide (74 g, 0.134 moles),
25 obtained as described in example 10, and 3,5-dichloro-pyridin-4-carbaldehyde (23.6 g, 0.134 moles) in CH₂Cl₂ (500 ml) under N₂ by controlling the temperature with a water-bath. The mixture was kept under stirring overnight, then cooled and treated with 5% HCl. The phases were separated and the acid one was re-extracted with CH₂Cl₂, washed with water/NaCl,
30 decoloured with charcoal, dried and concentrated under high vacuum. There were obtained

- 15 -

85.4 g of a crude which was used as such in the subsequent step. A sample of the crude was purified by flash chromatography (eluent: hexane/ethyl acetate 1:1).

¹H-NMR (CDCl₃): 8.60 (s, 2H); 7.77-6.68 (m, 4H); 3.80 (s, 3H).

5

Example 12

4-(3,5-Dichloro-pyridin-4-ylmethyl)-7-methoxy-2H-phthalazin-1-one

Hydrazine (18.4 ml, 0.378 moles) was added to a suspension of 3-(3,5-dichloro-pyridin-4-ylmethyl)-6-methoxy-3H-isobenzofuran-1-one (24.4 g, 0.126 moles), obtained as described in example 11, in CH₃OH (200 ml), under N₂. The mixture was heated under reflux for 1 hour, then kept overnight at room temperature, cooled in ice, and the solid was filtered, washed with very cold CH₃OH and dried in oven at 50°C under vacuum, to give 33.3 g of the title compound (yield: 80%). m.p.: 259-262°C.

¹H-NMR (CDCl₃): 12.34 (s, 1H); 8.64 (s, 2H); 8.19-7.54 (m 3H); 4.58 (s, 2H); 3.95 (s, 3H).

Example 13

15

4-Chloro-1-(3,5-dichloro-pyridin-4-ylmethyl)-6-methoxy-phthalazine

POCl₃ (22.2 ml, 230 mmoles) was added to a suspension of 4-(3,5-dichloro-pyridin-4-ylmethyl)-7-methoxy-2H-phthalazin-1-one (10 g, 25.5 mmoles), obtained as described in example 12, in acetonitrile (300 ml) and the mixture was heated under reflux. After 3 hours the solution was concentrated, taken up with CH₂Cl₂, with water, and the pH was brought to 7-8 with Na₂CO₃. The organic phases were decoloured with charcoal, dried and concentrated to give 10 g of the title compound (stoichiometric yield).

¹H-NMR (CDCl₃): 8.50 (s, 2H); 8.13-7.54 (m, 3H); 4.88 (s, 2H); 4.04 (s, 3H).

Example 14

[4-(3,5-Dichloro-pyridin-4-ylmethyl)-7-methoxy-phthalazin-1-yl]-hydrazine

25

Hydrazine hydrate (0.81 ml, 0.84 g, 16.8 mmoles) was added to a solution under N₂ of 4-chloro-1-(3,5-dichloro-pyridin-4-ylmethyl)-6-methoxy-phthalazine (2 g, 5.6 mmoles), obtained as described in example 13, in ethanol (30 ml), and the mixture was kept under reflux for 24 hours, then cooled in ice and the resultant precipitate was filtered, washed with ethanol and ethyl ether and dried under vacuum at 50°C to give 2.14 g of the title compound (stoichiometric yield). m.p.: 297-299°C.

30

- 16 -

¹H-NMR (DMSO/D₂O): 8.61 (s, 2H); 7.95-7.20 (m, 3H); 4.48 (s, 2H); 3.87 (s, 3H).

Example 15

6-(3,5-Dichloro-pyridin-4-ylmethyl)-9-methoxy-3-methyl-[1,2,4]-triazole-[3,4-a]-

5 phthalazine (Compound 2)

Acetic anhydride (0.2 ml, 0.22 g, 2.2 mmoles) was added to a suspension under N₂ of [4-(3,5-dichloro-pyridin-4-ylmethyl)-7-methoxy-phthalazin-1-yl]-hydrazine (0.7 g, 2 mmoles), obtained as described in example 14, in acetic acid, and the mixture was kept under reflux for 20 hours, then brought to small volume, taken up with CH₂Cl₂ and washed twice with
10 2.5% NaOH, then with water. The mixture was decoloured with charcoal, filtered over celite and concentrated under vacuum to give a solid which was triturated in ethyl ether to give 0.57 g of the title compound (yield: 77%). m.p.: 241.6-243.6°C.

¹H-NMR (CDCl₃): 8.55 (s, 2H); 8.06-7.34 (m, 3H); 4.76 (s, 2H); 4.03 (s, 3H); 2.46 (s, 3H).

Example 16

15 6-(3,5-Dichloro-pyridin-4-ylmethyl)-9-methoxy-tetrazolo[5,1-a]-phthalazine (Compound 3)

NaN₃ was added to a solution under N₂ of 4-chloro-1-(3,5-dichloro-pyridin-4-ylmethyl)-6-methoxy-phthalazine (1 g, 2.82 mmoles), obtained as described in example 13, in anhydrous DMF (20 ml) and the mixture was heated at 80°C overnight, then at 120°C for 7 hours, then poured into water (10 volumes) and extracted 3 times with CH₂Cl₂, dried and concentrated
20 under vacuum to give a solid which was purified by flash chromatography (eluent: 60:80 petrolatum/ethyl acetate 6:4). The eluate was crystallised from acetonitrile (75 ml) to give 0.36 g of the title compound (yield: 78.5%). m. p.: 275-276°C.

¹H-NMR (CDCl₃): 8.57 (s, 2H); 8.22 (d, 1H, JHH=8.7Hz); 8.246 (d, 1H, JHH=2.5Hz); 7.57 (dd, 1H); 4.89 (s, 2H); 4.10 (s, 3H).

25

Example 17

5-(3,5-Dichloro-pyridin-4-ylmethyl)-8-methoxy-1,3,3a,4-tetraaza-cyclopentan[a]naphthalene
(Compound 4)

NaH (0.14 g, 3.38 mmoles) was added to a solution under N₂ of 1H-tetrazole (0.315g, 4.5 mmoles) in anhydrous DMF (10 ml) and the mixture was kept under stirring for 2 hours. 4-
30 Chloro-1-(3,5-dichloro-pyridin-4-ylmethyl)-6-methoxy-phthalazine (0.8 g, 2.25 mmoles),

- 17 -

obtained as described in example 13, was added and it was heated at 80°C. then at 100°C overnight. The mixture was poured into water and extracted with CH₂Cl₂. The organic phase containing an insoluble was concentrated under vacuum and taken up with CH₃OH. A solid was obtained which, triturated with hot CH₃OH then cooled, was removed by filtration. The mother liquors were brought to dryness to give a solid which was flash chromatographed (eluent: CH₂Cl₂/CH₃OH 98:2), to yield a solid which, triturated with ethyl ether, gave 0.18 g of the title compound (yield: 22%). m.p.: 231.3-232.3°C (dec.).

¹H-NMR (CDCl₃): 8.77 (s, 1H); 8.56 (s, 2H); 8.09-8.03 (m, 2H); 7.40 (dd, 1H, JHH=9Hz, J2HH=2.7Hz); 4.76 (s, 2H); 4.04 (s, 3H).

Example 18

Evaluation of the PDE 4 enzyme inhibition

a) Purification of human polymorphonucleate leukocytes

The polymorphonucleate leukocytes (PMNs) were isolated from peripheral blood of healthy volunteers according to what described by Boyum A., Scand. J. Immunol., 1976, 5th suppl., 9). Shortly, the isolation of the PMNs was effected by Ficoll-Paque gradient centrifugation followed by sedimentation on dextrane and the erythrocyte contamination was eliminated by hypotonic lysis.

b) PDE 4 enzyme purification

The human PMNs were re-suspended in TRIS/HCl buffer (10mM pH 7.8) containing MgCl₂ (5mM), EGTA (4mM), mercaptoethanol (5mM), TRITON-X100 (1%), pepstatin A (1μM), PMSF (100μM) and leupeptin (1μM), and homogenised by Polytron. The homogenate was centrifuged at 25,000 x g for 30 minutes at 4°C and the supernatant was used for the PDE 4 enzyme purification by ion exchange chromatography using the FPLC technique according to what described by Schudt C. et al., Naunyn-Schmidberg's Arch. Pharmacol., 1991, 334, 682. The supernatant was seeded on an UNO Q12 column (Bio-Rad) and the enzyme was eluted by sodium acetate gradient from 50mM to 1M. The fractions containing enzymatic activity were collected, dialysed against water and concentrated. The resulting PDE 4 enzyme was stored at -20°C in the presence of ethylenglycole (30%, v/v) until the use.


- 18 -

c) PDE 4 enzyme inhibition

The enzyme activity was evaluated with an Amersham kit based on the SPA (Scintillation Proximity Assay) technique. The enzymatic reaction was effected in a total volume of 100 μ l of TRIS/HCl buffer (50mM, pH7.5), $MgCl_2$ (8.3mM), EGTA (1.7mM), cAMP (1 μ M) and [3H]cAMP (~100.000 dpm) as tracer. The compounds of the invention were added at the selected concentrations. The reaction was started by adding the enzyme (15 μ g protein/ml), went on for 40 minutes at 30°C and stopped by adding 50 μ l of suspension of SPA particles. The radioactivity due to the particles was measured in a β -emitting counter. The results are expressed as percent activity versus the control present in each experiment. The IC_{50} values were calculated over 9 equidistant concentrations in logarithmic scale using a 4-parameters logistic function by software. The compounds of the present invention showed to selectively inhibit PDE 4: for example. Compound 2 gave a value of IC_{50} =207nM.

- 19 -

1) Compounds of formula



wherein

A is a 5-7 membered heterocycle containing from 1 to 4 nitrogen atoms, optionally partially or totally unsaturated, and optionally substituted by a (C₁₋₄)alkyl group in turn optionally substituted;

10

Z is NH, methylene, a C₂₋₆ alkylene chain optionally branched and/or unsaturated and/or interrupted by a C₅₋₇ cycloalkyl residue;

Cy is phenyl or heterocycle optionally substituted by one or more substituents, or a COR₄ group wherein R₄ is hydroxy, alkoxy, amino optionally substituted by one or two (C₁₋₆)alkyl groups or by hydroxy;

15

R is a (C₁₋₆)alkyl or polyfluoro(C₁₋₆)alkyl group;

R₁ is hydrogen; a (C₁₋₈)-alkyl, (C₂₋₈)-alkenyl or (C₂₋₈)-alkynyl group optionally substituted by hydroxy, oxo, aryl or heterocycle, and optionally interrupted by one or more heteroatoms or heterogroups; a (C₁₋₄)alkoxy group or a (C₄₋₇)cycloalkoxy group optionally containing an oxygen atom and optionally substituted by a polar substituent in the cyclic moiety, aryloxy, aryl-(C₁₋₁₀)-alkoxy;

20

the N→O derivatives of the compounds of formula I and the pharmaceutically acceptable salts thereof.

2) Compounds according to claim 1 wherein Z is methylene or a C₂₋₆ alkylene chain.


25 3) Compounds according to claim 1 wherein Z is methylene or a C₂₋₆ alkylene chain; and Cy is a heterocycle optionally substituted by one or more substituents.

4) Compounds according to claim 1 wherein Z is methylene; and Cy is pyridine substituted by two substituents.

5) Process for the preparation of a compound according to claim 1 wherein Z is different from NH, characterised in that an acid of formula


30

- 20 -




5

wherein R and R₁ are as defined in claim 1, by reaction with formaldehyde/HCl gives a compound of formula




10

wherein R and R₁ are as above defined, which is oxidised and hydrolysed to give a compound of formula




15

wherein R and R₁ are as above defined, which is treated with a hydrohalogenidric acid and triphenylphosphine to give a compound of formula



20

25 wherein R and R₁ are as above defined, this compound being also obtainable from compound III by radicalic halogenation to give the compound of formula




30

- 21 -


wherein R and R₁ are as above defined, and X is chlorine or bromine, which gives compound V by treatment with triphenylphosphine; said compound of formula V treated with an aldehyde of formula




wherein Cy is as defined in claim 1 and Z'' is a C₁₋₅ alkylene chain optionally branched and/or unsaturated and/or interrupted by a C₅₋₇ cycloalkyl residue, or it is absent. in the presence of an organic base gives a compound of formula



wherein R, R₁, Z'' and Cy are as above defined, which is treated with hydrazine to give a compound of formula



wherein R, R₁ and Cy are as above defined and Z has the meanings reported in claim 1 but amino, which is treated with a halogenating agent to give a compound of formula



wherein R, R₁ and Cy are as above defined, and X is chlorine or bromine, which is treated with a suitable nucleophile or with hydrazine and then with a suitable acylating agent.

- 22 -

6. A pharmaceutical composition containing a therapeutically effective amount of a compound according to claim 1 in admixture with a suitable carrier.
7. A pharmaceutical composition according to claim 6 for the treatment of allergic and
5 inflammatory pathologies.
8. A pharmaceutical composition according to claim 6 for the treatment of respiratory diseases.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 99/07304

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D487/04 A61K31/5025 //(C07D487/04, 237:00; 257:00),
(C07D487/04, 237:00, 249:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	WO 99 32456 A (ALVAREZ BUILLA JULIO G ;GARCIA NAVIO JOSE LUIS (ES); SIRO HERRERO) 1 July 1999 (1999-07-01) the whole document	1-8
Y	WO 98 21208 A (BYK GULDEN LOMBERG CHEM FAB ;FLOCKERZI DIETER (DE)) 22 May 1998 (1998-05-22) cited in the application the whole document	1-8
Y	WO 93 07146 A (SYNTEX INC) 15 April 1993 (1993-04-15) the whole document	1-8
Y	EP 0 722 936 A (EISAI CO LTD) 24 July 1996 (1996-07-24) the whole document	1-8
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

4 February 2000

Date of mailing of the international search report

16/02/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Stellmach, J

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 99/07304

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 98 07430 A (KABASAWA YASUHIRO ; WATANABE NOBUHISA (JP); ABE SHINYA (JP); ADACHI) 26 February 1998 (1998-02-26) the whole document	1-8
Y, P	& EP 0 920 868 A (EISAI CO. LTD.) 9 June 1999 (1999-06-09) the whole document	1-8
Y	--- DAL PIAZ, V. ET AL.: "Novel Heterocyclic-Fused Pyridazinones as Potent and Selective Phosphodiesterase IV Inhibitors " J.MED.CHEM., vol. 40, no. 10, 1997, pages 1417-1421, XP002101978 WASHINGTON the whole document	1-8
Y	--- WO 98 04559 A (GUIBLIN ALEXANDER RICHARD ; MERCK SHARP & DOHME (GB); MOORE KEVIN W) 5 February 1998 (1998-02-05) the whole document	1-8
Y	--- DE 196 17 862 A (SCHERING AG) 30 October 1997 (1997-10-30) cited in the application the whole document	1-8
Y	--- EP 0 620 224 A (TAKEDA CHEMICAL INDUSTRIES LTD) 19 October 1994 (1994-10-19) cited in the application the whole document	1-8
Y	--- EP 0 548 923 A (TAKEDA CHEMICAL INDUSTRIES LTD) 30 June 1993 (1993-06-30) cited in the application the whole document -----	1-8

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/EP 99/07304

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9932456	A	01-07-1999	IT MI972806 A AU 2613299 A	21-06-1999 12-07-1999
WO 9821208	A	22-05-1998	AU 5317098 A EP 0937074 A NO 992282 A US 6008215 A	03-06-1998 25-08-1999 11-05-1999 28-12-1999
WO 9307146	A	15-04-1993	AT 183745 T AU 670544 B AU 2781592 A CA 2117059 A DE 69229874 D DE 69229874 T EP 0612321 A ES 2105920 A FI 941567 A HU 66969 A HU 9500113 A IL 103388 A JP 7500321 T MX 9205794 A NO 941210 A NZ 244660 A PT 100938 A, B US 5716954 A ZA 9207755 A	15-09-1999 25-07-1996 03-05-1993 15-04-1993 30-09-1999 09-12-1999 31-08-1994 16-10-1997 06-04-1994 30-01-1995 28-06-1995 30-09-1997 12-01-1995 01-04-1993 05-04-1994 26-05-1995 29-10-1993 10-02-1998 08-04-1994
EP 0722936	A	24-07-1996	AU 705229 B AU 3191995 A FI 961510 A NO 961397 A NZ 290952 A US 5849741 A CA 2173493 A CN 1135210 A HU 76067 A WO 9605176 A JP 8225541 A	20-05-1999 07-03-1996 29-05-1996 06-06-1996 27-05-1998 15-12-1998 22-02-1996 06-11-1996 30-06-1997 22-02-1996 03-09-1996
WO 9807430	A	26-02-1998	AU 3784897 A CA 2258079 A CN 1225586 A EP 0920868 A JP 10114657 A NO 990805 A	06-03-1998 26-02-1998 11-08-1999 09-06-1999 06-05-1998 16-04-1999
WO 9804559	A	05-02-1998	AU 3551997 A AU 3553997 A BG 103112 A CA 2260774 A CZ 9900181 A EP 0915877 A EP 0915875 A WO 9804560 A NO 990304 A PL 331072 A ZA 9706591 A	20-02-1998 20-02-1998 30-09-1999 05-02-1998 16-06-1999 19-05-1999 19-05-1999 05-02-1998 25-03-1999 21-06-1999 18-08-1998

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/EP 99/07304

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9804559 A		HR 980035 A	30-06-1999
DE 19617862 A	30-10-1997	NONE	
EP 0620224 A	19-10-1994	AT 168376 T	15-08-1998
		CA 2120997 A	13-10-1994
		DE 69411616 D	20-08-1998
		DE 69411616 T	19-11-1998
		JP 6345767 A	20-12-1994
		US 5492909 A	20-02-1996
EP 0548923 A	30-06-1993	AT 168113 T	15-07-1998
		CA 2086164 A	28-06-1993
		DE 69226157 D	13-08-1998
		DE 69226157 T	22-10-1998
		JP 6116272 A	26-04-1994
		US 5369104 A	29-11-1994
		JP 5271233 A	19-10-1993

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

This Page Blank (uspto)